

This surgery is a very safe, painless procedure performed in a hospital or outpatient surgery center, and it generally takes less than 20 minutes to complete. There are no needles and no stitches involved. The eye is numbed with anesthetic eye drops, and you will receive either an IV or oral sedation to help you relax.

Surgery Date: _____ **Arrival Time:** _____ **Location:** _____

Do NOT eat or drink anything after midnight the night before your surgery.

The Day of Your Procedure:

- You **CAN** take medication for high blood pressure or heart medication the morning of your surgery with a small sip of water. (Only drink enough to get the medicine down.)
- If you are **Diabetic**, do **NOT** take your medication (pill or insulin) the morning of your surgery. Bring your medication with you, and you may take it immediately after surgery.
- Wear comfortable clothes.
- Bring an adult to drive you home. You will receive either an IV or oral sedation, and you will need to be driven home.
- Be sure to have all eye drop prescriptions filled before surgery, and bring the medication with you when you report to the surgery center.

Your first follow up appointment is at: _____

on: _____ **time:** _____

- Patient acknowledges receipt of bag, sunglasses, tape, Bromday, Besivance, and Durezol.**

Patient Signature

Date

**If you have any questions Mon-Fri 8AM-5PM, please call your surgery coordinator,
Tina Doty at 901-255-5616**

Getting Ready for Surgery:

Drop 1:

BROMDAY

Fill Prescription. (Start using on _____) Eye: Right Left

This is a non-steroid drop to decrease inflammation. Use one drop in the operative eye(s) once a day. If using more than one drop, remember to wait five minutes before using your next eye drop. Do NOT discontinue use until instructed by your doctor.

Drop 2:

DUREZOL

Fill Prescription. (Start using on _____) Eye: Right Left

This is a steroid drop to decrease inflammation. Use one drop in the operative eye(s) four times a day.

If using more than one drop, remember to wait 5 minutes before using the next drop. DO NOT discontinue use until instructed by your doctor.

You will continue it at home four times a day.

IF YOU FAIL TO GET THIS MEDICATION, YOU WILL NEED TO RECEIVE A STEROID INJECTION IN YOUR EYE IN ORDER TO CONTROL INFLAMMATION.

Drop 3:

BESIVANCE

Fill Prescription. (Start using on _____) Eye: Right Left

This is an antibiotic drop used to prevent infection after surgery.

Use one drop in the operative eye(s) three times a day. If using more than one drop, remember to wait five minutes before using your next eye drop. Do NOT discontinue use until instructed by your doctor.