Pre-Operative Instructions

Surgery Date & Time:



Acular [Start using on _____

This is a non-steroid drop to decrease inflammation. Use one drop in the operative eye(s) four times a day beginning four days before surgery. We recommend scheduling your drops with breakfast, lunch, dinner, and bedtime. Remember to wait five minutes before using your next eye drop. Do NOT discontinue use until instructed by your doctor.



Zymar [Start using on ____]

This is an antibiotic drop used to prevent infection after surgery. Use one drop in the operative eye(s) four times a day beginning one day before surgery. We recommend scheduling your drops with breakfast, lunch, dinner, and bedtime. Remember to wait five minutes before using your next eye drop. Do NOT discontinue use until instructed by your doctor.



Pred Forte [Start using on ____]

This is a steroid solution used to decrease inflammation and regulate healing. Use one drop in the operative eye(s) four times a day beginning on the day of surgery. Remember to shake the bottle gently before each use and to wait five minutes before using another eye drop. Do NOT discontinue use until instructed by your doctor.

Please note: You may need to continue to use these drops for more than one week. This will be determined at your one week post-op appointment. The drops may sting a little, but it is imperative that you continue your drops as directed.

On the Day of Surgery

- Do NOT eat or drink anything after midnight the morning of your surgery.
- You CAN take medication for high blood pressure or heart medication the morning of your surgery with a small sip of <u>water</u>. Only drink enough to get the medicine down.
- Wear comfortable clothes. Do not wear your hair in a ponytail or braid, and do not wear any makeup.
- You must have a responsible adult accompany you on the day of your surgery and for your follow-up appointment the next day.
- Bring your black surgery kit with you to surgery.
- Be sure to have your **Zymar** prescription filled before surgery, and bring the medication with you when you report to the surgery center.

